REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden. to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave bla	nk) 2. REPORT DATE 3/3/99	3. REPORT TYPE AND DATES final	COVERED
4. TITLE AND SUBTITLE		le rint	DING NUMBERS
	ect effects of die	sel fuel on	
Direct and indirect effects of diesel fuel on microphytobenthos and meiofauna in saltmarsh			014-96-1-0242
sediments			
	est for the section of the section o		
6. AUTHOR(S)	ni and Kevin @arman		
Inomas S. Planer	ii and kevin earman		
7. PERFORMING ORGANIZATION N			ORMING ORGANIZATION ORT NUMBER
Dept. of EEO Bio			
310 Dinwiddie Ha		1	
Tulane Universit		·	
New Orleans 701	118		
			MEANINE / MANUTARINE
	GENCY NAME(S) AND ADDRESS(ES) Research, Code 114		NSORING/MONITORING NCY REPORT NUMBER
		100	
800 N. Quincy St			
Arlington, VA 2	22217		
11. SUPPLEMENTARY NOTES		internation and activities activiti	
11. SUPPLEMENTARY NOTES			
i			
12a. DISTRIBUTION / AVAILABILITY	STATEMENT	N - C 1 D(C	TOIRIITION CODE
distribution uni	limited		
distribution uni	limited		
distribution uni	limited		
distribution un	limited	19990315	
13. ABSTRACT (Maximum 200 wor	rds)	19990315	5 090
13. ABSTRACT (Maximum 200 wor Resultsprovide	os) several insights in	1999031	090 of diesel
13. ABSTRACT (Maximum 200 wor Resultsprovide s contaminants on	os) several insights in benthic food webs.	1999031	of diesel
13. ABSTRACT (Maximum 200 wor Resultsprovide a contaminants on the base of this	os) several insights in benthic food webs. s food web. Ths co	1999031	of diesel gae constitute lts from 1994
13. ABSTRACT (Maximum 200 work Resultsprovide a contaminants on the base of this and 1995 experis	os) several insights in benthic food webs. s food web. Ths co ments indicates the	1999031 Ito the influence of Benthic microals Insistency of result In robustness of our	of diesel gae constitute lts from 1994 c observations
13. ABSTRACT (Maximum 200 work Resultsprovide contaminants on the base of this and 1995 experiments of reduced coper	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh	1999031 Ito the influence of Benthic microals Insistency of result In robustness of our lanced algal biomas	of diesel gae constitute tts from 1994 c observations
13. ABSTRACT (Maximum 200 work Resultsprovide contaminants on the base of this and 1995 expering of reduced coperations of high presence of high	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel	1999031	of diesel gae constitute tts from 1994 cobservations in the ile the broad
13. ABSTRACT (Maximum 200 work Resultsprovide a contaminants on the base of this and 1995 experiments of reduced coperations of the presence of high responses of the contaminants on the contaminants on the contaminants of the	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w	1999031 Ito the influence of Benthic microals ensistency of result erobustness of our lanced algal biomas contamination. While were similar, differ	of diesel gae constitute Its from 1994 r observations ss in the ile the broad erences in the
13. ABSTRACT (Maximum 200 work Resultsprovide of contaminants on the base of this and 1995 experiments of reduced coperations of the presence of high responses of the response of the compares the contaminant of the compares of the compare	os) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi	1999031 Ito the influence of Benthic microals ensistency of result erobustness of our lanced algal biomas contamination. While were similar, differences are consistent	of diesel gae constitute Its from 1994 c observations ss in the ile the broad erences in the int with the
13. ABSTRACT (Maximum 200 work Resultsprovide of contaminants on the base of this and 1995 experiments of reduced coperate of presence of high responses of the hypothesis that	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi the MS community i	1999031 Ito the influence of Benthic microals Insistency of result Insistency of res	of diesel gae constitute tts from 1994 cobservations in the ile the broad erences in the at with the to contaminants.
13. ABSTRACT (Maximum 200 work Resultsprovide of the base of this and 1995 experiments of reduced coperates of the responses of the hypothesis that Microalgal bioms	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi the MS community i ass in sediments is	dto the influence of Benthic microals on sistency of result anced algal biomas contamination. While ties are consistents more sensitive is controlled primar	of diesel gae constitute ts from 1994 cobservations in the ile the broaderences in the at with the to contaminants.
13. ABSTRACT (Maximum 200 work Resultsprovide contaminants on the base of this and 1995 experiments of reduced coperates of high responses of the hypothesis that Microalgal biomeiofaunal graz	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi the MS community i ass in sediments is ing pressure. Mort	1999031 Ito the influence of Benthic microal onsistency of result anced algal biomas contamination. White the sare consistent is more sensitive to controlled primar ality to meiofaunce	of diesel gae constitute Its from 1994 r observations ss in the ile the broad erences in the nt with the to contaminants. rily by al grazers
13. ABSTRACT (Maximum 200 work Resultsprovide contaminants on the base of this and 1995 experiments of reduced coperates of high responses of the hypothesis that Microalgal biomomeiofaunal graz releases algae	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi the MS community i ass in sediments is ing pressure. Mort from grazing pressu	1999031 to the influence of Benthic microal on sistency of result anced algal biomas contamination. White ties are consistent are consistent as more sensitive to controlled primary ality to meiofaunce. The algae groups	of diesel gae constitute ts from 1994 cobservations in the ile the broad erences in the ht with the to contaminants. Tily by al grazers ow until they
13. ABSTRACT (Maximum 200 work Resultsprovide contaminants on the base of this and 1995 experiments of reduced coperates of high responses of the hypothesis that Microalgal biomomeiofaunal graz releases algae become N-limite	several insights in benthic food webs. s food web. Ths coments indicates the pod grazing and enh levels of diesele two communities we meiofaunal communithe MS community is ing pressure. Mort from grazing pressud. Enhanced ammoni	1999031. Ito the influence of Benthic microals insistency of result anced algal biomas contamination. White ties are consistent is more sensitive to controlled primary ality to meiofaunce and the algae group production in contamination.	of diesel gae constitute Its from 1994 r observations ss in the ile the broad erences in the nt with the to contaminants. rily by al grazers ow until they diesel-
13. ABSTRACT (Maximum 200 work Resultsprovide a contaminants on the base of this and 1995 experiments of reduced coperate of high responses of the response of the hypothesis that Microalgal biomomeiofaunal graz releases algae become N-limite contaminated se	ds) several insights in benthic food webs. s food web. Ths coments indicates the pod grazing and enh levels of diesel e two communities we meiofaunal communithe MS community i ass in sediments is ing pressure. Mort from grazing pressud. Enhanced ammonidiments then allows	1999031 Ito the influence of Benthic microals of result in the robustness of our lanced algal biomas contamination. White ties are consistent in the secont of the second	of diesel gae constitute Its from 1994 c observations is in the ile the broad erences in the int with the to contaminants. rily by al grazers ow until they diesel- develop futther.
13. ABSTRACT (Maximum 200 work Resultsprovide a contaminants on the base of this and 1995 experiments of reduced coperate presence of high responses of the hypothesis that Microalgal biomomeiofaunal graz releases algae become N-limite contaminated segments	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi the MS community i ass in sediments is ing pressure. Mort from grazing pressu d. Enhanced ammoni diments then allows	1999031 Ito the influence of Benthic microals on sistency of result anced algal biomas contamination. White the sare consistent is more sensitive to some sensitive to some sensitive to ality to meiofaunce. The algae group production in the salgal blooms to some sensitive to salgal blooms to sa	of diesel gae constitute Its from 1994 r observations ss in the ile the broad erences in the nt with the to contaminants. rily by al grazers ow until they diesel- develop futther.
13. ABSTRACT (Maximum 200 work Resultsprovide a contaminants on the base of this and 1995 experiments of reduced coperate presence of high responses of the hypothesis that Microalgal biomomeiofaunal graz releases algae become N-limite contaminated segments	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi the MS community i ass in sediments is ing pressure. Mort from grazing pressu d. Enhanced ammoni diments then allows	1999031 Ito the influence of Benthic microals on sistency of result anced algal biomas contamination. White the sare consistent is more sensitive to some sensitive to some sensitive to ality to meiofaunce. The algae group production in the salgal blooms to some sensitive to salgal blooms to sa	of diesel gae constitute Its from 1994 r observations ss in the ile the broad erences in the nt with the to contaminants. rily by al grazers ow until they diesel- develop futther.
13. ABSTRACT (Maximum 200 work Resultsprovide of contaminants on the base of this and 1995 experiments of reduced coperate of responses of the response of the hypothesis that Microalgal biomomeiofaunal graz releases algae become N-limite contaminated setting, our experiments.	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi the MS community i ass in sediments is ing pressure. Mort from grazing pressu d. Enhanced ammoni diments then allows iments have allowed h blooms in diesel-	1999031 Ito the influence of Benthic microals on sistency of result anced algal biomas contamination. White the sare consistent is more sensitive to some sensitive to some sensitive to ality to meiofaunce. The algae group production in the salgal blooms to some sensitive to salgal blooms to sa	of diesel gae constitute Its from 1994 r observations ss in the ile the broad erences in the nt with the to contaminants. rily by al grazers ow until they diesel- develop further. he general r.
13. ABSTRACT (Maximum 200 work Resultsprovide a contaminants on the base of this and 1995 experiments of reduced coperate presence of high responses of the hypothesis that Microalgal biomomeiofaunal graz releases algae become N-limite contaminated segments	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi the MS community i ass in sediments is ing pressure. Mort from grazing pressu d. Enhanced ammoni diments then allows iments have allowed h blooms in diesel-	1999031 Ito the influence of Benthic microals on sistency of result anced algal biomas contamination. White the sare consistent is more sensitive to some sensitive to some sensitive to ality to meiofaunce. The algae group production in the salgal blooms to some sensitive to salgal blooms to sa	of diesel gae constitute Its from 1994 r observations ss in the ile the broad erences in the nt with the to contaminants. rily by al grazers ow until they diesel- develop futther. he general r. 15. NUMBER OF PAGES 3
13. ABSTRACT (Maximum 200 work Resultsprovide of contaminants on the base of this and 1995 experiments of reduced coperate of responses of the response of the hypothesis that Microalgal biomomeiofaunal graz releases algae become N-limite contaminated setting, our experiments.	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi the MS community i ass in sediments is ing pressure. Mort from grazing pressu d. Enhanced ammoni diments then allows iments have allowed h blooms in diesel-	1999031 Ito the influence of Benthic microals on sistency of result anced algal biomas contamination. White the sare consistent is more sensitive to some sensitive to some sensitive to ality to meiofaunce. The algae group production in the salgal blooms to some sensitive to salgal blooms to sa	of diesel gae constitute Its from 1994 r observations ss in the ile the broad erences in the nt with the to contaminants. rily by al grazers ow until they diesel- develop further. he general r.
13. ABSTRACT (Maximum 200 work Resultsprovide of contaminants on the base of this and 1995 experiments of reduced coperate of responses of the response of the hypothesis that Microalgal biomomeiofaunal graz releases algae become N-limite contaminated setting, our experiments.	ds) several insights in benthic food webs. s food web. Ths co ments indicates the pod grazing and enh h levels of diesel e two communities w meiofaunal communi the MS community i ass in sediments is ing pressure. Mort from grazing pressu d. Enhanced ammoni diments then allows iments have allowed h blooms in diesel-	1999031 Ito the influence of Benthic microals on sistency of result anced algal biomas contamination. White the sare consistent is more sensitive to some sensitive to some sensitive to ality to meiofaunce. The algae group production in the salgal blooms to some sensitive to salgal blooms to sa	of diesel gae constitute Its from 1994 r observations ss in the ile the broad erences in the nt with the to contaminants. rily by al grazers ow until they diesel- develop futther. he general r. 15. NUMBER OF PAGES 3

FINAL REPORT

GRANT #: 96PR01926-00

PRINCIPAL INVESTIGATOR: Thomas S. Bianchi

INSTITUTION: Tulane University

GRANT TITLE: Direct and indirect effects of diesel fuel on microphytobenthos and meiofauna in saltmarsh

sediments

AWARD PERIOD: - 30 Oct. 1998

OBJECTIVES: To determine if(i) algal blooms in contaminated sediments are the result of direct or indirect stimulation from diesel or from a reduction in meiofaunal grazing; (ii) mortality to copepods in contaminated sediments is from direct toxicity from diesel, anoxic conditions, or a combination of the two; and (iii) Cletocamptus deitersi (a meiofaunal copepod) thrives in contaminated sediments because of reduced competition from other species, or exploitation of altered algal resources.

APPROACH: Microcosms experiments were performed to determine the effects of diesel-contaminated sediment on microalgal-meiofaunal interactions. Natural sediments were defaunated, then manipulated to control meiofaunal abundance and species composition, as well as diesel contamination. Treatments were compared to determine if reduced grazing or altered nutrient availability in diesel-contaminated sediments lead to algal blooms. Toxicity tests were used to determine interspecific differences in sensitivity to diesel and oxygen depletion. The influence of diesel on meiofaunal competitive interactions (grazing rates) was measured in communities manipulated to mimic those that occur in diesel-contaminated sediments. Microcosm experiments were performed to determine if enhanced N availability in diesel-contaminated sediments was from decaying carcasses of meiofauna, or altered sedimentary biogeochemical processes. HPLC analysis was employed to determine changes in sedimentary microphytobenthos, and how those changes relate to copepod feeding.

ACCOMPLISHMENTS: The effects of altered grazing and nutrient availability were examined in two microcosm experiments. In the first, we determined if diesel contamination influences NH_4^+ flux from sediments. Results indicate an enhanced flux of NH_4^+ from contaminated sediments. NH_4^+ concentrations remained very low in contaminated sediments incubated in the light, because microalgae consumed it as fast as it was produced. Addition of NH_4^+ to uncontaminated sediments did not stimulate algal growth, however, indicating that N availability was not the primary factor limiting algal growth.

In a second experiment, we examined the hypothesis that reduced grazing led to the formation of algal blooms. Meiofaunal abundance was manipulated to remove copepods and other larger grazers from sediment. Algal responses to this manipulation of grazing were compared to those that occurred in diesel-contaminated sediments. Both removal of grazers and contamination with diesel produced a 2-fold increase in algal biomass over a 4 days. Microscopic and HPLC analyses indicate that the increased algal biomass was almost entirely from growth of diatoms.

Experiments were also conducted to distinguish between the effects of hypoxia vs direct toxicity of diesel on meiofaunal copepods. In all species examined, we observed that hypoxia greatly increased the toxicity of diesel-contaminated sediment. *Cletocamptus deitersi*, a species that seems to be resistant to diesel toxicity, was the least sensitive of the species tested.

In previous microcosm experiments using natural communities, we observed that nematodes and C. deitersi thrive and exhibit enhanced grazing rates on microphytobenthos, leading to the hypothesis that C. deitersi and nematodes benefit from a release from competition for food when other copepods are killed. This hypothesis was tested by experimentally manipulating C. deitersi and nematode abundances to simulate conditions that occur in diesel-contaminated sediment. Grazing by C. deitersi and nematodes increased in the absence of copepod competitors. Among nematodes, enhanced grazing on algae was limited to a diatom feeder, Ptycholaimellus sp. Our evidence indicates that increased grazing by C. deitersi in contaminated sediments is partially from reduced competition, and partially a response to the enhanced algal biomass in contaminated sediments.

Our research indicates that algal blooms in diesel-contaminated sediments are a consequence of reduced grazing (from meiofaunal mortality) over short periods (5-14 days), followed by a second growth phase that is fueled by enhanced N fluxes. We performed two experiments to examine the source of this excess nitrogen. The first examined the possibility that N comes from decaying carcasses of animals killed via diesel toxicity. Results indicate that nitrogen released from decaying carcasses occurs over very short periods (3-4 days), well before algae become N-limited. Thus, excess N that fuels growth of algae must come from other N pools in the sediment.

Observations above led to an experiment (performed last fall) in which we examined in more detail the biogeochemistry of N dynamics in diesel-contaminated sediment. We are in the final stages of analyzing samples from this experiment. We are examining the hypothesis that altered microbial activity in diesel-contaminated sediment may result in increased metabolism of refractory marsh detritus, which contains geoplymers that may have "complexed" DIN; this could result in the release of organic and inorganic nitrogen. We are measuring total hydrolyzable and dissolved free amino acids (THAA) in experimental sediments from diesel-contaminated and control sediments, as well as N and P dynamics in inorganic pools (PO4, NH4, NO3, and NO2), and bulk sediment nitrogen. Analysis of inorganic pools has been completed, and analysis of amino acids and bulk sediment is underway. Preliminary analysis indicates that diesel contamination dramatically enhances the NH₄ (as previously documented) and PO₄ flux from contaminated sediments relative to controls. Evidence of enhanced PO₄ flux is new, and may contribute to the formation of microphytobenthos blooms in contaminated sediments. When amino-acid and bulk-sediment analyses are completed, we should have a much clearer picture of the origin of excess nitrogen that fluxes from contaminated sediments.

CONCLUSIONS: Results provide several insights into the influence of diesel contaminants on benthic food webs. Benthic microalgae constitute the base of this food web, and thus their dramatic increase in response to contaminants is ecologically significant. The consistency of results from 1994 and 1995 experiments indicates the robustness of our observations of reduced copepod grazing and enhanced algal biomass in the presence of high levels of diesel contamination. While the broad responses of the two communities were similar, differences in the response of the meiofaunal communities are consistent with the hypothesis that the MS community is more sensitive to contaminants. Microalgal biomass in sediments is controlled primarily by meiofaunal grazing pressure. Mortality to meiofaunal grazers (which occurs as a consequence of the interactive effects of hypoxia and diesel toxicity) releases algae from grazing pressure. The algae grow until they become N-limited. Enhanced NH₄+ production in diesel-contaminated sediments then allows algal blooms to develop further. Thus, our experiments have allowed us to identify the general process by which blooms occur in diesel-contaminated sediments.

SIGNIFICANCE: Collectively, our research has yielded an understanding of how and why benthic communities in salt marshes respond to diesel. Diesel contamination dramatically alters benthic invertebrate community structure, grazing on microphytobenthos, and nitrogen biogeochemistry. These cascading effects lead to algal blooms and fundamental changes in competitive interactions among benthic invertebrates.

AWARD INFORMATION: T. Bianchi was awarded tenure in 1998.

PUBLICATIONS AND ABSTRACTS

- Bennett, A., T.S., Bianchi, J.C. Means, & K. Carman. The effects of PAH contamination and grazing on the abundance and composition of microphytobenthos in salt marsh sediments (Pass Fourchon, LA): I. A microcosm experiment. J. Exp. Mar. Biol. Ecol. (Submitted; in revision)
- Carman, K.R., Fleeger, J.W., & S.M. Pomarico. Does chronic hydrocarbon contamination alter the response of benthic communities to diesel contamination? Mar. Environ. Res. (Submitted, in revision).
- Bennett, A., T.S. Bianchi, & J. C. Means. The effects of PAH contamination and grazing on the abundance and composition of microphytobenthos in salt marsh sediments (Pass Fourchon, LA):II. A field study. Estuar. Coastal Shelf Sci. (Submitted).
- Carman, K.R. & J.C. Means. Sediment bacterial response to hydrocarbon contamination in two coastal saltmarshes. Estuaries. (In prep)
- Carman, K.C., Bianchi, T.S., & J.W. Fleeger. Microalgal blooms in hydrocarbon-contaminated saltmarsh sediments: The influences of grazing and nitrogen availability. Mar. Ecol.

Prog. Ser. (In prep)

Bianchi, T.S. & K.R. Carman. The effects of PAH contamination on the availability of nitrogen to microphytobenthos in marsh sediments Org. Geochem. (In prep).

Abstracts

1998 Benthic Ecology Meeting (4)

Invited seminars given by Carman at Florida State University, Old Dominion Univ., Inst. for Marine Sciences (UNC Chapel Hill marine lab), Univ. North Carolina Chapel Hill, Univ. South Carolina

1997 American Society for Limnology & Oceanography (3)